FLOYD CHEROKEE MEDICAL CENTER POLICY AND PROCEDURE MANUAL PATIENT CARE SERVICES



TITLE: Interchange of Quinolones by a Clinical Pharmacist	Policy No.: C-PCS-06-029
Purpose: To establish a protocol for the interchange of quinolones according to the results of culture and sensitivity reports to assure the most sensitive quinolone is prescribed.	
Reference Standards: MM 05.01.01	·

Policy:

The clinical pharmacist may interchange quinolones according to the patient's culture and sensitivity results by writing a per policy order from the physician.

- 1. When the following criteria are met, the clinical pharmacist may interchange quinolones so that the most sensitive quinolone is administered to the patient.
 - ◆ The patient is currently receiving a quinolone.
 - ♦ The culture and sensitivity results show a difference in the sensitivities (minimal inhibitory concentration, i.e., MIC) for quinolones.
 - The patient is not currently receiving the most sensitive quinolone.
- 2. The clinical pharmacist may enter a per policy order from the physician to change the prescribed quinolone to the most sensitive quinolone.
- 3. The physician will sign the per policy order in the patient's chart within forty-eight hours.
- 4. A physician may enter "Dispense as Written" to override the quinolone interchange.

Reference Policy: PH 11-011